

Manifestations of Extra Dimensions in a Neutrino Telescope

A. Nicolaidis and D.T. Papadamou

Department of Theoretical Physics, University of Thessaloniki, GR-54006 Thessaloniki,
Greece.

Abstract

Theories with large extra dimensions provide the possibility that a flavor neutrino, localized in a 3+1 brane, can mix with a singlet neutrino living in the bulk. This mixing leads to unconventional patterns of neutrino matter oscillations and we examine in details how these oscillations depend upon two parameters: the brane-bulk coupling ξ and the effective mass μ of the flavor neutrino inside matter. We find that high energy ($E \geq 50$ GeV) ν_μ neutrinos, to be detected by neutrino telescopes, can give signals of extra dimensions. With a 1 km^3 neutrino telescope extra dimensions with radius down to $1\mu m$ can be tested directly, while for smaller radius an indirect evidence can be established.

Introduction. The standard model (SM) of strong and electroweak interactions has been extremely successful. It provides a consistent theoretical framework within which we can analyze and understand all available experimental data. Still we know that it cannot be regarded as the final theory. Any attempt to include quantum gravity leads to a unified theory where two disparate scales coexist, the electroweak scale ($M_W \sim 1$ TeV) and the Planck scale ($M_{Pl} \sim 10^{19}$ GeV). Quantum radiative corrections then, especially in the scalar sector (Higgs field) of the theory, tend to mix the scales and without an incredible amount of fine tuning, will always equalize them (hierarchy problem).

A novel approach has been suggested to alleviate the hierarchy problem [1]. Our four-dimensional world is embedded in a higher dimensional space with D dimensions ($D = 4 + n$). While the SM fields are constrained to live on the 4-dimensional brane, gravity can freely propagate in the higher-dimensional space (bulk). The fundamental scale M_f of gravity in D dimensions is related to the observed 4-dimensional Planck scale M_{Pl} by

$$M_{Pl}^2 = M_f^{2+n} V_n \quad (1)$$

where V_n is the volume of the extra space. For a torus configuration

$$V_n = (2\pi)^n R_1 R_2 \dots R_n \quad (2)$$

with R_i ($i = 1, 2, \dots, n$) the radii of extra dimensions. Then for a sufficiently large volume V_n the fundamental scale of gravity M_f can become as low as M_W . In this radical way the hierarchy problem ceases to exist as such.

Besides the graviton, a sterile neutrino can freely propagate in the extra dimensions. Since the extra dimensions are compact, a bulk neutrino will appear at the 4-dimensional brane as a Kaluza-Klein (KK) tower, i.e. an infinite number of 4-dimensional spinors. The Yukawa coupling of the standard lepton doublet, the Higgs scalar and the right-handed bulk neutrino will provide a mixing between the left handed neutrino of the standard model and the KK modes. In the next part we present this mixing and we obtain the eigenvectors and the eigenvalues of the mass matrix. In the third part we focus our attention into neutrino oscillations, inside matter. Compared to the usual oscillations, novel features appear since now we have a coupled system of infinite degrees of freedom. The pattern of oscillations depends upon two parameters : the coupling ξ between the left-handed neutrino and the KK states and the effective mass μ of the flavor neutrino inside matter. We study in detail this dependence. In the final part we suggest that high energy cosmic ν_μ neutrinos can probe more efficiently the bulk of the space. These neutrinos, after traversing the Earth, will be detected at future neutrino telescopes. We present detailed spectra of the cosmic ν_μ as function of the energy and the incident angle. At the end we present our conclusions .

The Brane-Bulk Model. The available gravity experiments imply that any deviation from the Newton law should occur at distances r less than 1 mm. Then, to satisfy equations (1)-(2), we need more than two extra dimensions. To simplify our investigation we consider a single large extra dimension, assuming all other dimensions smaller and not affecting our analysis.

There is already an extensive literature on neutrinos living in extra dimensions [2]-[13]. We start by considering the action for a massless bulk fermion Ψ living in 5 dimensions

$$S_\Psi = \int d^4x dy \bar{\Psi} \Gamma^\mu \partial_\mu \Psi \quad M = 0, 1, \dots, 5 \quad (3)$$

Ψ is decomposed as

$$\Psi = \begin{pmatrix} N_R \\ N_L \end{pmatrix} \quad (4)$$

each component admitting a Fourier expansion

$$N(x, y) = \frac{1}{\sqrt{2\pi R}} N_0(x) + \sum_{n=1}^{\infty} \frac{1}{\sqrt{\pi R}} \left(N_n(x) \cos\left(\frac{ny}{R}\right) + \hat{N}_n(x) \sin\left(\frac{ny}{R}\right) \right) \quad (5)$$

Thus at the brane the bulk fermion appears as a KK tower. Since the momentum in the extra compact dimension is quantized, each KK mode appears as having a mass n/R ($n = 1, 2, \dots$). Consider now the coupling of the standard left-handed lepton doublet to the bulk neutrino

$$\frac{h}{\sqrt{M_f}} \bar{L} H N_R \delta(y) \quad (6)$$

where H is the Higgs scalar doublet and h a dimensionless Yukawa coupling. After the Higgs field develops a vacuum expectation value (v), we get the mass terms

$$m \bar{\nu}_L (N_{R0} + \sqrt{2} \sum_{n=1}^{\infty} N_{Rn}) \quad (7)$$

with $m \sim h v M_f / M_{Pl}$. Notice that for $M_f \sim 1 TeV$, $h \sim 1$, we obtain $m \sim 10^{-4} eV$ [2]-[7]. Also, the left handed zero mode N_{L0} decouples from the spectrum and remains massless. Denoting

$$\Psi_R = \begin{pmatrix} N_{R0} \\ N_{Ri} \end{pmatrix} \quad \Psi_L = \begin{pmatrix} \nu_L \\ N_{Li} \end{pmatrix} \quad i = 1, 2, \dots \quad (8)$$

the mass term is $\bar{\Psi}_L M \Psi_R$ with the mass matrix M given by [4]

$$M = \begin{pmatrix} m & \sqrt{2}m & \sqrt{2}m & \cdots \\ 0 & 1/R & 0 & \cdots \\ 0 & 0 & 2/R & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} \quad (9)$$

The evolution of the neutrino states is determined by the Hamiltonian

$$H = \frac{1}{2E_\nu} M M^T \quad (10)$$

In terms of rescaled variables, $H = X/(2E_\nu R^2)$, where the dimensionless matrix X becomes (we keep k rows and columns)

$$X = \begin{pmatrix} (2k+1)\xi^2 & \sqrt{2}\xi & 2\sqrt{2}\xi & 3\sqrt{2}\xi & \cdots & (k-1)\sqrt{2}\xi \\ \sqrt{2}\xi & 1 & 0 & 0 & \cdots & 0 \\ 2\sqrt{2}\xi & 0 & 4 & 0 & \cdots & 0 \\ 3\sqrt{2}\xi & 0 & 0 & 9 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ (k-1)\sqrt{2}\xi & 0 & 0 & 0 & 0 & (k-1)^2 \end{pmatrix} \quad (11)$$

with $\xi = mR$. For a neutrino propagating inside matter the upper-left element is replaced $(2k+1)\xi^2 \rightarrow (2k+1)\xi^2 + \mu$, where μ is proportional to the effective mass of the neutrino inside matter. For a ν_μ neutrino $\mu = \sqrt{2}E_\nu R^2 G_F N_n$ with N_n the neutron density. The eigenvalues λ_n^2 of the matrix X satisfy the equation

$$\left[\mu - \lambda^2 + \xi^2(\lambda\pi) \cot(\lambda\pi) \right] \prod_{n=1}^{\infty} (n^2 - \lambda^2) = 0 \quad (12)$$

The corresponding eigenvector to λ_n^2 is $B_n = (e_{n0}, e_{n1}, e_{n2}, \dots)$ with [4]

$$e_{nk} = -\frac{k\sqrt{2}\xi}{(k^2 - \lambda_n^2)} e_{n0} \quad k = 1, 2, 3, \dots \quad (13)$$

$$e_{n0}^2 \left(\frac{1}{2} + \frac{\pi^2 \xi^2}{2} + \frac{\mu}{2\lambda_n^2} + \frac{(\lambda_n - \frac{\mu}{\lambda_n})^2}{2\xi^2} \right) = 1 \quad (14)$$

Neutrino Mixing. Before presenting the numerical solution to equation (12) it is useful to mention some qualitative features

i) small ξ

For $\xi = 0$ there is no coupling between the SM neutrino and the KK states. The left-handed neutrino remains massless, $\lambda_0 = 0$, while for a neutrino propagation inside matter $\lambda_0 = \sqrt{\mu}$. The other eigenvalues are $\lambda_n = n$ ($n = 1, 2, 3, \dots$). As ξ starts to rise $\lambda_n \simeq n + \xi^2/(n\pi)$ (for large n)

ii) large ξ

For large ξ , λ_n tend to $n + 1/2$. An expansion for large ξ and large n provides

$$\lambda_n \simeq (n + \frac{1}{2})(1 - \frac{1}{\xi^2 \pi^2}) \quad (15)$$

iii) resonance structure

The amount of admixture of the n -th eigenstate B_n to the neutrino state ν is determined by $|e_{n0}|^2$ and it is becoming maximal when the resonance condition

$$\mu_R = \lambda_n^2 \quad (16)$$

holds (see equation 14). Since $\lambda_n \simeq n$, the resonance condition is simplified to $\mu \simeq n^2$ ($n = 1, 2, 3, \dots$). For small ξ values the resonance is narrow, while for large ξ values the resonance is broad and more than one state might be involved. Recall that for a neutral medium

$$\mu = \frac{G_F R^2 \rho E_\nu}{\sqrt{2} M_N} \quad (17)$$

with ρ the density of the medium and M_N the nucleon mass. The requirement $\mu \simeq 1$ (strong oscillation phenomena) leads to a relationship between the model parameters (R), the medium properties (ρ) and the neutrino energy (E_ν)

$$\left(\frac{\rho}{10 \frac{gr}{cm^3}}\right) \left(\frac{E_\nu}{100 GeV}\right) \left(\frac{R}{1 \mu m}\right)^2 \simeq 1 \quad (18)$$

iv) Level repulsion

When there is no coupling among the SM neutrino and the KK states ($\xi = 0$), the eigenvalues cross each other (see dashed lines in figure 1). When the off diagonal elements become non-zero ($\xi \neq 0$), the well-known phenomenon of level repulsion appears : as the lower eigenvalue increases by increasing μ , by approaching the resonance region ($\mu \simeq 1$) levels off and reaches asymptotically the value 1, thus avoiding crossing the next eigenvalue. Similarly the higher eigenvalue starts at $\lambda \simeq 1$, increases as μ increases and when encounters the second resonance ($\mu \simeq 4$) levels off at the value 2. (see the figure 1)

v) graphical solution

The eigenvalues λ_n^2 can be found graphically as the intersection points of the two curves (see the figure 2)

$$y_1 = \lambda^2 - \mu \quad (19)$$

$$y_2 = \xi^2(\lambda\pi) \cot(\lambda\pi) \quad (20)$$

Most of the qualitative figures described above can be seen by inspecting the figure 2.

Oscillating Neutrinos and Neutrino Telescopes. From the previous analysis it is clear that for neutrinos traveling inside the Earth, strong oscillation phenomena occur whenever the condition (18) is satisfied. Thus for $R \sim 1\mu m$, we need neutrinos of energy $E \sim 100$ GeV, while a smaller radius is probed by even higher energy neutrinos. It is clear that neutrino telescopes [14] which are able to detect ν_μ neutrinos from ~ 20 GeV up to ~ 50 TeV are best suited for this study. At this early stage of investigation we adopt a simple model where the ν_μ flavor state mixes only with a bulk neutrino. We expect this simple model will indicate the salient features of the more complicated physical phenomenon.

Once a ν_μ flavor state is produced, its time evolution will be governed by the equation

$$|\nu(t)\rangle = \sum_n e_{n0}^* \exp\left(\frac{-i\lambda_n^2 t}{2E_\nu R^2}\right) |B_n\rangle \quad (21)$$

where the summation over n is carried up to $n_{max} = [E_\nu R]$ The amplitude for the transition $\nu_\mu \rightarrow \nu_\mu$ is

$$A = \sum_n |e_{n0}|^2 \exp\left(\frac{-i\lambda_n^2 t}{2E_\nu R^2}\right) \quad (22)$$

Figure 3 shows the probability $P(\nu_\mu \rightarrow \nu_\mu)$ for a neutrino traversing a medium of constant density, as a function of distance. The value we used for the density of the medium corresponds to the density of the Earth's mantle. We used also $R = 1\mu m$. A clear dip develops at the appropriate length scale. The probability $P(\nu_\mu \rightarrow \nu_\mu)$ as a function of energy is presented in figure 4. Notice that an impressive resonance structure appears at the energy which corresponds to $\mu \simeq 1$ and all ν_μ disappear. At the resonance, reading the value of the energy and using equation (18) we can extract the value of the radius R .

It is useful to evaluate the transition probability for a neutrino traversing the Earth. For the Earth's density we adopt a model where the core is a sphere with constant density 11 gr/cm^3 while the mantle, surrounding the core has constant density 4.4 gr/cm^3 [15]. We consider a neutrino which for time t_1 propagates inside the mantle, for time t_2 crosses the core and it takes time t_3 within the mantle again. The amplitude $A(\nu_\mu \rightarrow \nu_\mu)$ is given by

$$A = \sum_{n,l,r} e_{n0}^* \exp\left(\frac{-i\lambda_n^2 t_1}{2E_\nu R^2}\right) D_{nl} \exp\left(\frac{-i\tilde{\lambda}_l^2 t_2}{2E_\nu R^2}\right) D_{lr}^{-1} \exp\left(\frac{-i\lambda_r^2 t_3}{2E_\nu R^2}\right) e_{r0} \quad (23)$$

where tilded (untilded) parameters correspond to core (mantle) and D is the rotation matrix from one set of eigenvectors to the other

$$D_{nl} = \sum_k e_{nk} \tilde{e}_{lk}^* \quad (24)$$

The probability $P(\nu_\mu \rightarrow \nu_\mu)$ for a neutrino going through the center of the Earth, as a function of the energy is shown in figure 5. The resonance dip is now displaced.

How efficiently can a neutrino telescope unravel the underlying dynamics? Without entering into technical details and specifications of a neutrino telescope, we can recall that by determining the direction of the produced muon, we also fix the direction of the parent muon neutrino. Thus we can have a zenith coverage of incident neutrino from $\theta = 0$ (neutrinos going through the center of Earth) to $\theta \simeq \pi/2$ (neutrinos just scratching the Earth, before being detected). In the absence of any oscillations the neutrino flux is θ -independent. If oscillations are present, by varying θ we vary the distance x traveled by ν_μ (x varies continuously from 10 km for $\theta = \pi/2$ to 13000 km for $\theta = 0$). In our case the $\theta \simeq \pi/2$ neutrino flux remains undistorted by matter effects and represents therefore the initial incident neutrino flux. The deduced ratio $I(\theta)/I(\theta = \pi/2)$ ($I(\theta)$ is the inferred intensity of neutrino flux at angle θ) reflects then the probability for a muon neutrino to oscillate. The energy of the incident neutrino is determined for the contained events [16]. Thus we have a set of events, where both the distance x and the energy of the neutrino E_ν are determined, allowing to extract the theoretical parameters (ξ and R). Neutrino induced events with energy up to 300 – 400 GeV are expected to be contained within a 1 km^3 volume and therefore we deduce that 1 km^3 neutrino telescope can provide direct evidence for an extra dimension with $R \geq 1\mu\text{m}$. For smaller R , we expect the resonance structure to shift to higher energies and thus the resonance dip will not be directly seen. Still we may have an indirect evidence for strong oscillations by examining the ratio $r = E_s/E_p$, with $E_s(E_p)$ the number of stopping (passing) muons [17]. If at $\theta = 0$ the number of highly energetic ν_μ ($E_\nu \geq 400 \text{ GeV}$) is depleted, then we expect the ratio r to increase for $\theta = 0$ compared to the same ratio for $\theta \simeq \pi/2$.

Another important scale in the neutrino oscillations is the resonance oscillation length L_R . A significant depletion of ν_μ occurs whenever the distance traveled x is equal to the oscillation length L , defined by

$$L = 2\pi \frac{E_\nu R^2}{\lambda_1^2 - \lambda_0^2} \quad (25)$$

At the resonance, $\mu \simeq 1$, L becomes

$$L_R = \frac{2\sqrt{2} \pi M_N}{G_F \rho (\lambda_1^2 - \lambda_0^2)} \quad (26)$$

Whenever both equations (18) and (26) are satisfied, we observe the maximum depletion of ν_μ ($P(\nu_\mu \rightarrow \nu_\mu) \simeq 0$). The eigenvalues difference $\lambda_1^2 - \lambda_0^2$ depends upon ξ . For small ξ values the resonance is narrow and the difference $\lambda_1^2 - \lambda_0^2$ is small. The corresponding L_R is big, much bigger than the Earth's diameter. For example for $\xi = 0.05$ and with $\rho = 10 \text{ gr/cm}^3$ we find $L_R = 23571$ km. In that case strong oscillations cannot be observed. For $\xi = 2.2$ (the value we opted), we find $L_R = 6000$ km, smaller than Earth's size, and strong oscillations are visible.

Conclusions. We analyzed the situation where the left-handed neutrino ν_μ of the SM mixes with a right handed neutrino experiencing a large extra space dimension. This mixing induces neutrino oscillations, even if the neutrino itself is massless. We encountered novel features compared to the standard oscillations, since the mixing involves an infinity of KK states. A rich resonance structure appears accompanied by many spikes. We presented in detail the first resonance ($\mu \simeq 1$). A similar study can be realized for the second resonance ($\mu \simeq 4$). We emphasized that 1 km^3 neutrino telescope may be used as an instrument to explore the bulk space and establish unambiguously the existence of extra dimensions with $R \geq 1\mu\text{m}$. We didn't address the issues related to low energy neutrino data (solar neutrinos, atmospheric neutrinos), since for the R values we used, there is no significant mixing of low energy neutrinos with the bulk states. After all, the recent neutrino data (SNO [18]) disfavor the presence of a sterile neutrino. Altogether our analysis shows that neutrino mixing in models with large extra dimensions is an interesting phenomenon and deserves further study.

References

- [1] I. Antoniadis Phys. Lett. **B246** (1990) 377; N. Arkani-Hamed, S. Dimopoulos and G. Dvali , Phys. Lett. **B429** (1998) 263; I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali , Phys. Lett. **B436** (1998) 257.
- [2] N. Arkani-Hamed, S. Dimopoulos, G.Dvali and J. March-Russell hep-ph/9811448.
- [3] K. Dienes, E. Dudas and T. Ghergetta Nucl. Phys. **B557** (1999) 25 .
- [4] R. Barbieri, P. Creminelli and A. Strumia, Nucl. Phys. **B585** (2000) 28.
- [5] R. Mohaparta, S. Nandi and Perez-Lorenzana, Phys. Lett. **B466** (1999) 115.
- [6] A. Lukas, P. Ramond, A. Romanino and G Ross, Phys. Lett. **B495** (2000) 136.
- [7] G. Dvali and A. Smirnov, Nucl. Phys. **B563** (1999) 63.
- [8] K. Dienes and I Sarcevic, Phys. Lett. **B500** (2001) 133.
- [9] A. Das and O. Kong, Phys. Lett. **B470** (1999) 149.
- [10] D. Caldwell, R. Mohaparta and S. Yellin hep-ph/0102279.
- [11] N. Cosme, J. Frre, Y. Gouverneur, F. Ling, D. Monderen and V. Van Elewyck, Phys. Rev. **D63** (2001) 113018.
- [12] C.S. Lam and J.N Ng hep-ph/0104129.
- [13] A.Gouvea, G. Giudice, A. Strumia, K. Tobe, hep-ph/0107156.
- [14] The AMANDA Collaboration Astropart, Phys **13**, 1 (2000) .
The NESTOR Collaboration Nucl. Phys. **Proc.Suppl 87** (2000) 448.
The ANTARES Collaboration Nucl. Phys. **Proc.Suppl 81** (2000) 174.
- [15] A. Nicolaidis Phys. Lett. **B200** (1988) 553.
- [16] Ivone F.M. Albuquerque and George F. Smoot hep-ph/0102078.
- [17] P.Lipari, M.Lusignoli and F. Sartogo, Phys. Rev. Lett. **74** (1995) 4384;
P.Lipari, M.Lusignoli Phys. Rev. **D57** (1998) 3842
- [18] Q. Ahmad et al., SNO Collaboration, nucl-ex/0106015.




Figure 1: Eigenvalues λ as a function of μ . Dashed lines for $\xi = 0$ (no coupling among the SM neutrino and the KK states) and the solid lines for $\xi = 0.2$




Figure 2: Dash line for $f(\lambda) = \lambda^2 - \mu$ and solid line for $f(\lambda) = \xi^2(\lambda\pi) \cot(\lambda\pi)$ with $\xi = 0.2$




Figure 3: The probability $P(\nu_\mu \rightarrow \nu_\mu)$ for a neutrino of energy 100GeV traversing a medium with density $\rho = 4.4 \text{ gr/cm}^3$ (close to the density of the Earth's mantle) as a function of distance.




Figure 4: The probability $P(\nu_\mu \rightarrow \nu_\mu)$ for a neutrino traversing 10649 Km through the mantle of Earth (with density $\rho = 4.4 \text{ gr/cm}^3$), as a function of energy.




Figure 5: The probability $P(\nu_\mu \rightarrow \nu_\mu)$ for a neutrino going through the center of the Earth , as a function of energy.